

# SYMMETRIC AND ASYMMETRIC HYDRODYNAMICS FOR THE FACILITATED EXCLUSION PROCESS VIA MAPPING

BASED ON J.W. WITH O. BLONDEL, M. SASADA, M. SIMON AND L. ZHAO

Clément Erignoux, INRIA Lille

*Interacting Particle Systems and Hydrodynamic Limits*  
*Montréal, March 24 2022*

# SYMMETRIC SIMPLE EXCLUSION PROCESS (SSEP) ON $\mathbb{Z}$

---

- ▷ Configuration  $\eta \in \Omega := \{0, 1\}^{\mathbb{Z}}$ , with  $\eta_x = 1$  for an occupied site,  $\eta_x = 0$  for an empty site.
- ▷ **Stirring dynamics**: two neighboring sites are exchanged at rate 1.
- ▷ Initial profile  $\rho_0 : \mathbb{R} \rightarrow [0, 1]$  fixed, **initial configuration** e.g.  $\eta_x(0) = 1$  w.p.  $\rho_0(x/N)$ .

Then, the **empirical measure** on a diffusive timescale

$$\pi_{tN^2}^N = \frac{1}{N} \sum_{x \in \mathbb{Z}} \eta_x(tN^2) \delta_{x/N}$$


converges in a weak sense to  $\rho(t, u) du$ , where  $\rho$  is the **solution to the heat equation**

$$\begin{cases} \partial_t \rho = \partial_{uu} \rho \\ \rho(0, \cdot) = \rho_0 \end{cases} .$$

# FACILITATED EXCLUSION PROCESS (FEP)

---

Similar to [Gonçalves, Landim, Toninelli '08], but with stronger kinetic constraint



Markov generator  $\mathcal{L}f(\eta) = \sum_{x \in \mathbb{Z}} c_{x,x+1}(\eta) \{f(\eta^{x,x+1}) - f(\eta)\},$

with

$$c_{x,x+1}(\eta) = p\eta_{x-1}\eta_x(1 - \eta_{x+1}) + (1 - p)\eta_{x+2}\eta_{x+1}(1 - \eta_x).$$


The parameter  $p \in [0, 1]$  tunes the asymmetry, and  $\eta^{x,x+1}$  is the configuration where sites  $x$  and  $x + 1$  have been exchanged.

- ▷ Bernoulli product measures are **not stationary**.
- ▷ **No mobile cluster** to mix the configuration (cooperative model).

# FACILITATED EXCLUSION PROCESS (FEP)

---

Similar to [Gonçalves, Landim, Toninelli '08], but with stronger kinetic constraint



Markov generator  $\mathcal{L}f(\eta) = \sum_{x \in \mathbb{Z}} c_{x,x+1}(\eta) \{f(\eta^{x,x+1}) - f(\eta)\},$

with

$$c_{x,x+1}(\eta) = p\eta_{x-1}\eta_x(1 - \eta_{x+1}) + (1 - p)\eta_{x+2}\eta_{x+1}(1 - \eta_x).$$


The parameter  $p \in [0, 1]$  tunes the asymmetry, and  $\eta^{x,x+1}$  is the configuration where sites  $x$  and  $x + 1$  have been exchanged.

- ▷ Bernoulli product measures are **not stationary**.
- ▷ **No mobile cluster** to mix the configuration (cooperative model).

# FACILITATED EXCLUSION PROCESS (FEP)

---

Similar to [Gonçalves, Landim, Toninelli '08], but with stronger kinetic constraint



Markov generator  $\mathcal{L}f(\eta) = \sum_{x \in \mathbb{Z}} c_{x,x+1}(\eta) \{f(\eta^{x,x+1}) - f(\eta)\},$

with

$$c_{x,x+1}(\eta) = p\eta_{x-1}\eta_x(1 - \eta_{x+1}) + (1 - p)\eta_{x+2}\eta_{x+1}(1 - \eta_x).$$

The parameter  $p \in [0, 1]$  tunes the asymmetry, and  $\eta^{x,x+1}$  is the configuration where sites  $x$  and  $x + 1$  have been exchanged.

- ▷ Bernoulli product measures are **not stationary**.
- ▷ **No mobile cluster** to mix the configuration (cooperative model).

# HYDRODYNAMIC LIMIT FOR THE SYMMETRIC FEP

Theorem (Blondel, E', Simon, Sasada 2018 & BES 2021)

Given  $\rho_0$ , consider the **symmetric** ( $p = 1 - p = 1/2$ ) process  $\eta(t)$  on  $\mathbb{T}_N := \{0, 1, \dots, N\}$  started from

$$\mu^N = \mu_0^N := \bigotimes_{x \in \mathbb{T}_N} \text{Ber}(\rho_0(x/N)).$$

For any smooth compactly supported  $H$

$$\frac{1}{N} \sum_{x \in \mathbb{T}_N} H(x/N) \eta_x(tN^2) \xrightarrow[N \rightarrow \infty]{\mathbb{P}} \int_{[0,1]} H(u) \rho(t, u) du$$

where  $\rho$  is solution to the parabolic Stefan problem  $\rho(0, u) = \rho_0(u)$  and

$$\boxed{\rho_0 > 1/2}$$

$$\boxed{\rho_0 \in [0, 1]}$$

$$\partial_t \rho = \frac{1}{2} \partial_{uu} \left\{ \frac{2\rho - 1}{\rho} \right\}$$

$$\partial_t \rho = \frac{1}{2} \partial_{uu} \left\{ \frac{2\rho - 1}{\rho} \mathbf{1}_{\{\rho \geq 1/2\}} \right\}.$$

# HYDRODYNAMIC LIMIT FOR THE SYMMETRIC FEP

Theorem (Blondel, E', Simon, Sasada 2018 & BES 2021)

Given  $\rho_0$ , consider the **symmetric** ( $p = 1 - p = 1/2$ ) process  $\eta(t)$  on  $\mathbb{T}_N := \{0, 1, \dots, N\}$  started from

$$\mu^N = \mu_0^N := \bigotimes_{x \in \mathbb{T}_N} \text{Ber}(\rho_0(x/N)).$$

For any smooth compactly supported  $H$

$$\frac{1}{N} \sum_{x \in \mathbb{T}_N} H(x/N) \eta_x(tN^2) \xrightarrow[N \rightarrow \infty]{\mathbb{P}} \int_{[0,1]} H(u) \rho(t, u) du$$

where  $\rho$  is solution to the parabolic Stefan problem  $\rho(0, u) = \rho_0(u)$  and

$$\boxed{\rho_0 > 1/2}$$

$$\boxed{\rho_0 \in [0, 1]}$$

$$\partial_t \rho = \frac{1}{2} \partial_{uu} \left\{ \frac{2\rho - 1}{\rho} \right\}$$

$$\partial_t \rho = \frac{1}{2} \partial_{uu} \left\{ \frac{2\rho - 1}{\rho} \mathbf{1}_{\{\rho \geq 1/2\}} \right\}.$$

# TYPES OF CONFIGURATIONS

---

Four types of configurations, depending on the **critical density**  $\rho_c = 1/2$ .

- **Low density** : if  $\rho < 1/2$

Frozen configurations

$$\mathcal{F} = \{\eta \in \Omega \mid \eta_x \eta_{x+1} \equiv 0\}$$



Transient Bad configurations

$$\mathcal{TB} = \{\eta \in \Omega \mid \eta_x \eta_{x+1} \neq 0\}.$$



- **Large density** : if  $\rho > 1/2$ ,

Ergodic configurations

$$\mathcal{E} = \{\eta \in \Omega \mid (1-\eta_x)(1-\eta_{x+1}) \equiv 0\}$$



Transient Good configurations

$$\mathcal{TG} = \{\eta \in \Omega \mid (1-\eta_x)(1-\eta_{x+1}) \neq 0\}$$





# GRAND CANONICAL MEASURES

---

Because of kinetic constraint, Bernoulli product measures are not stationary for the dynamics. Canonical measures can be defined as **uniform measures on the ergodic components** with fixed number of particles.

The symmetric FEP is actually **reversible w.r.t. a family of explicit supercritical distributions**  $\pi_\rho$ , for  $\rho \in (1/2, 1]$ .

- ▷  $\pi_\rho$  is supported on the **infinite ergodic component**.
- ▷  $\pi_\rho$  is a Bernoulli product measure conditioned to having isolated empty sites (ergodic component)
- ▷  $\pi_\rho$  exhibits **long-range correlations** as  $\rho \searrow 1/2$ .

# ENTROPY TOOLS AND EQUILIBRIUM DISTRIBUTIONS

---

The most classical techniques for hydrodynamic limits are based on **entropy bounds** between the measure  $\mu_t^N$  of the process at time  $t$  and its reference measures  $\pi_\alpha$ , namely

▷ Guo, Papanicolaou and Varadhan's **entropy method**,

$$H(\mu_t^N \mid \pi_\rho) \leq CN,$$

▷ Yau's **relative entropy method**

$$H(\mu_t^N \mid \pi_{\rho_t}) = o(N).$$

**Supercritical case**, in the transient regime,  $\mu_t^N$  is not supported on ergodic configurations, whereas the grand canonical measures  $\pi_\rho$  are  $\Rightarrow$  entropy estimate fails. In particular, we need to prove that the ergodic component is reached before the diffusive timescale  $\tau = O(N^2)$ .

**General case**, no real hope of using entropy methods : no reference measures because the two phase's stationary states have **disjoint supports**, and no smooth solutions to the hydrodynamic limit.

# STRATEGY OF PROOF

---

- ▷ **Supercritical case**, GPV's entropy method can be adapted, by proving that the ergodic component is reached in a subdiffusive time.
  
- ▷ **General case:**
  - ▶ entropy methods cannot be used, so we adapt Funaki's scheme for parabolic Stephan problems.
  - ▶ The **one-block estimate** is based on a De Finetti-type decomposition for translation invariant stationary states.
  - ▶ The two blocks estimate is bypassed by directly proving that the **Young measure is a dirac**.

# HYDRODYNAMIC LIMIT FOR THE ASYMMETRIC FEP

## Theorem (E', Simon, Zhao 2022)

Given  $\rho_0$ , consider the **asymmetric** ( $p \in (1/2, 1]$ ) process  $\eta(t)$  started from

$$\mu^N = \mu_0^N := \bigotimes_{x \in \mathbb{Z}} \text{Ber}(\rho_0(x/N)).$$

For any smooth compactly supported  $H$

$$\frac{1}{N} \sum_{x \in \mathbb{Z}} H(x/N) \eta_x(tN) \xrightarrow[N \rightarrow \infty]{\mathbb{P}} \int_{\mathbb{R}} H(u) \rho(t, u) du$$

where  $\rho$  is the **unique entropy solution** to the hyperbolic Stefan problem

$$\begin{cases} \partial_t \rho + (2p - 1) \partial_u \left\{ \mathfrak{H}(\rho) \mathbf{1}_{\{\rho \geq 1/2\}} \right\} \\ \rho(0, u) = \rho_0(u) \end{cases}, \quad \text{where} \quad \mathfrak{H}(\rho) = \frac{(1 - \rho)(2\rho - 1)}{\rho}.$$

# HYDRODYNAMIC LIMIT FOR THE ASYMMETRIC FEP

## Theorem (E', Simon, Zhao 2022)

Given  $\rho_0$ , consider the **asymmetric** ( $p \in (1/2, 1]$ ) process  $\eta(t)$  started from

$$\mu^N = \mu_0^N := \bigotimes_{x \in \mathbb{Z}} \text{Ber}(\rho_0(x/N)).$$

For any smooth compactly supported  $H$

$$\frac{1}{N} \sum_{x \in \mathbb{Z}} H(x/N) \eta_x(tN) \xrightarrow[N \rightarrow \infty]{\mathbb{P}} \int_{\mathbb{R}} H(u) \rho(t, u) du$$

where  $\rho$  is the **unique entropy solution** to the hyperbolic Stefan problem



$$\begin{cases} \partial_t \rho + (2p - 1) \partial_u \left\{ \mathfrak{H}(\rho) \mathbf{1}_{\{\rho \geq 1/2\}} \right\} \\ \rho(0, u) = \rho_0(u) \end{cases}, \quad \text{where} \quad \mathfrak{H}(\rho) = \frac{(1 - \rho)(2\rho - 1)}{\rho}.$$

# Possible strategies of proof

- ▷ GPV's **entropy method** for hyperbolic systems ? No two-blocks estimate in the asymmetric case.
- ▷ Yau's **relative entropy method** ? Only useful until the first shock, and even so, not at all straightforward for two-phased systems, and no smooth solution a priori even before the shock because of the Stefan problem.
- ▷ Fritz's **compensated compactness** arguments ? Blackbox tools, very technical, and requires adding up some lower-order stirring dynamics.
- ▷ **Attractiveness** ? A priori not available here.


# MAPPING WITH A FACILITATED ZR PROCESS

---



# MAPPING WITH A FACILITATED ZR PROCESS


---






# MAPPING WITH A FACILITATED ZR PROCESS

---




# MAPPING WITH A FACILITATED ZR PROCESS

---




# MAPPING WITH A FACILITATED ZR PROCESS

---





# MAPPING WITH FZRP : DYNAMICS

---




# MAPPING WITH FZRP : DYNAMICS

---




# MAPPING WITH FZRP : DYNAMICS

---



# MAPPING WITH FZRP : DYNAMICS



⇒ If the exclusion process is driven by the facilitated generator, the corresponding **facilitated zero-range process (FZRP)** seen from the tagged empty site is driven by the generator

$$\mathcal{L}^{zr} g(\omega) = \sum_{y \in \mathbb{Z}} \mathbf{1}_{\{\omega_y \geq 2\}} \left\{ pg(\omega^{y,y+1}) + (1-p)g(\omega^{y,y-1}) - g(\omega) \right\}.$$

## PROPERTIES OF THE FZRP

---

- ▷ Since the function  $k \mapsto \mathbf{1}_{\{k \geq 2\}}$  is non-decreasing, this "facilitated" zero-range process is **attractive**: the evolution of two such processes  $\omega$  and  $\zeta$  can be coupled in such a way that

$$\omega(0) \leq \zeta(0) \quad \Rightarrow \quad \omega(t) \leq \zeta(t) \quad \forall t.$$

- ▷ The equilibrium/stationary distributions for the FZRP are product geometric measures **with no empty sites**, and density  $\alpha > 1$ , i.e. with marginals


$$\nu_\alpha(\omega_0 = k) = \mathbf{1}_{\{k \geq 1\}} \frac{1}{\alpha} \left(1 - \frac{1}{\alpha}\right)^{k-1}$$

- ▷ Even with attractiveness, coupling arguments are tricky, because the **process is not ergodic**: filling an empty site with a particle is irreversible for the FZRP, and equilibrium states only exist in the supercritical phase  $\alpha > 1$ .




# STRATEGY OF PROOF, HDL FOR THE FEP

---




# STRATEGY OF PROOF, HDL FOR THE FEP

---




# STRATEGY OF PROOF, HDL FOR THE FEP

---



# STRATEGY OF PROOF, HDL FOR THE FEP

---



# HYDRODYNAMICS FOR THE FZRP

## Theorem (E', Simon, Zhao 2022)

Given an initial profile  $\alpha_0$ , consider the **asymmetric** ( $p \in (1/2, 1]$ ) FZRP  $\omega(t)$ . Assuming that for any smooth compactly supported  $H$ , under the initial distribution,

$$\frac{1}{N} \sum_{y \in \mathbb{Z}} H(y/N) \omega_y \xrightarrow[N \rightarrow \infty]{\mathbb{P}} \int_{\mathbb{R}} H(v) \alpha_0(v) dv$$

then for any  $t > 0$

$$\frac{1}{N} \sum_{y \in \mathbb{Z}} H(y/N) \omega_y(tN) \xrightarrow[N \rightarrow \infty]{\mathbb{P}} \int_{\mathbb{R}} H(v) \alpha(t, v) dv$$

where  $\alpha$  is the **unique entropy solution** to the hyperbolic Stefan problem

$$\partial_t \alpha + (2p - 1) \partial_v \left\{ \frac{(\alpha - 1)}{\alpha} \mathbf{1}_{\{\alpha \geq 1\}} \right\} \quad \alpha(0, u) = \alpha_0(u).$$

↳ Hydrodynamic limit for attractive particle systems on  $\mathbb{Z}^d$ , F. Rezakhanlou.


# MACROSCOPIC MAPPING

- ▷ Denote  $X_0 = X_0(t)$  the position of the tagged empty site in the FEP, and  $\nu_t[\rho] = \lim_{N \rightarrow \infty} X_0(t)/N$  its macroscopic position at time  $t$ .
- ▷ The macroscopic position of the tagged empty site is formally written as

$$\nu_t[\alpha] = \nu_0 + \int_0^\infty \alpha_0(v) - \alpha(t, v) dv.$$

- ▷ Space variable  $y$  for  $\omega$  corresponding to  $x$  in  $\eta$ ? **Number of empty sites** between  $X_0$  and  $x$ . At the **macroscopic scale**  $u = x/N$ ,  $v = y/N$ , we can write

$$y = y(x) = \sum_{x'=X_0}^x (1 - \eta_{x'}) \Rightarrow v = v(u) = \int_{\nu_t}^u (1 - \rho(u')) du'$$




# MACROSCOPIC MAPPING

- ▷ Denote  $X_0 = X_0(t)$  the position of the tagged empty site in the FEP, and  $\nu_t[\rho] = \lim_{N \rightarrow \infty} X_0(t)/N$  its macroscopic position at time  $t$ .
- ▷ The macroscopic position of the tagged empty site is formally written as

$$\nu_t[\alpha] = \nu_0 + \int_0^\infty \alpha_0(v) - \alpha(t, v) dv.$$

- ▷ Space variable  $y$  for  $\omega$  corresponding to  $x$  in  $\eta$ ? **Number of empty sites** between  $X_0$  and  $x$ . At the **macroscopic scale**  $u = x/N$ ,  $v = y/N$ , we can write

$$y = y(x) = \sum_{x'=X_0}^x (1 - \eta_{x'}) \Rightarrow v = v(u) = \int_{\nu_t}^u (1 - \rho(u')) du'$$




# MACROSCOPIC MAPPING

- ▷ Denote  $X_0 = X_0(t)$  the position of the tagged empty site in the FEP, and  $\nu_t[\rho] = \lim_{N \rightarrow \infty} X_0(t)/N$  its macroscopic position at time  $t$ .
- ▷ The macroscopic position of the tagged empty site is formally written as

$$\nu_t[\alpha] = \nu_0 + \int_0^\infty \alpha_0(v) - \alpha(t, v) dv.$$

- ▷ Space variable  $y$  for  $\omega$  corresponding to  $x$  in  $\eta$ ? **Number of empty sites** between  $X_0$  and  $x$ . At the **macroscopic scale**  $u = x/N$ ,  $v = y/N$ , we can write

$$y = y(x) = \sum_{x'=X_0}^x (1 - \eta_{x'}) \Rightarrow v = v(u) = \int_{\nu_t}^u (1 - \rho(u')) du'$$






# MACROSCOPIC MAPPING

- ▷ Denote  $X_0 = X_0(t)$  the position of the tagged empty site in the FEP, and  $\nu_t[\rho] = \lim_{N \rightarrow \infty} X_0(t)/N$  its macroscopic position at time  $t$ .
- ▷ The macroscopic position of the tagged empty site is formally written as

$$\nu_t[\alpha] = \nu_0 + \int_0^\infty \alpha_0(v) - \alpha(t, v) dv.$$

- ▷ Space variable  $y$  for  $\omega$  corresponding to  $x$  in  $\eta$ ? **Number of empty sites** between  $X_0$  and  $x$ . At the **macroscopic scale**  $u = x/N$ ,  $v = y/N$ , we can write

$$y = y(x) = \sum_{x'=X_0}^x (1 - \eta_{x'}) \Rightarrow v = v(u) = \int_{\nu_t}^u (1 - \rho(u')) du'$$



# MAPPING HYDRODYNAMICS

---

Now, to prove the HDL for the FEP given that of the FZRP, one can use that

$$\frac{1}{N} \sum_{x \in \mathbb{Z}} \eta_x(tN^2) H(x/N) \simeq \frac{1}{N} \sum_{y \in \mathbb{Z}} \omega_y(tN^2) [H \circ u_t](y/N) + O(1/N),$$

where  $u = u_t(v)$  is the inverse mapping of  $v = v_t(u)$  seen earlier. Assuming everything is smooth, thanks to the HDL for the FZRP

$$\frac{1}{N} \sum_{y \in \mathbb{Z}} \omega_y(tN^2) [H \circ u_t](y/N) \simeq \int_{\mathbb{R}} \alpha(t, v) [H \circ u_t](v) dv,$$

and by a change of variable  $v \mapsto u_t(v)$ , the right hand side becomes

$$\int_{\mathbb{R}} \rho(t, u) H(u) du,$$

where  $\rho$  is given by  $\rho(u) = \frac{\alpha}{1+\alpha}(v)$ .

$\mapsto$  *Problem*: everything needs to be smoothed out, because of the hyperbolic equation, and because of the Stefan problem.

# MAPPING HYDRODYNAMICS

---

Now, to prove the HDL for the FEP given that of the FZRP, one can use that

$$\frac{1}{N} \sum_{x \in \mathbb{Z}} \eta_x(tN^2) H(x/N) \simeq \frac{1}{N} \sum_{y \in \mathbb{Z}} \omega_y(tN^2) [H \circ u_t](y/N) + O(1/N),$$

where  $u = u_t(v)$  is the inverse mapping of  $v = v_t(u)$  seen earlier. Assuming everything is smooth, thanks to the HDL for the FZRP

$$\frac{1}{N} \sum_{y \in \mathbb{Z}} \omega_y(tN^2) [H \circ u_t](y/N) \simeq \int_{\mathbb{R}} \alpha(t, v) [H \circ u_t](v) dv,$$

and by a change of variable  $v \mapsto u_t(v)$ , the right hand side becomes

$$\int_{\mathbb{R}} \rho(t, u) H(u) du,$$

where  $\rho$  is given by  $\rho(u) = \frac{\alpha}{1+\alpha}(v)$ .

$\mapsto$  *Problem*: everything needs to be smoothed out, because of the hyperbolic equation, and because of the Stefan problem.

## CURRENT WORK

---

- ▷ Phase transition(s) for the FEP/CLG in higher dimensions,  
*with A. Roget, A. Shapira and M. Simon.*
  
- ▷ Effect of boundary interactions on the FEP,  
*with M. Simon, PhD project.*

# THANKS FOR YOUR ATTENTION !

---

## RELATED WORKS - FEP

- ▷ E., Simon, Zhao, arxiv - 2202.04469 (2022).
- ▷ Blondel, E. and Simon, *Probability and Mathematical Physics* (2021).
- ▷ Blondel, E., Sasada and Simon, *An. de l'IHP - Prob. et Stat.* (2020).

## ATTRACTIVENESS & STEFAN PROBLEMS

- ▷ Seppäläinen. Translation invariant exclusion processes (2008).
- ▷ Kipnis, Landim, *Scaling Limits of Interacting Particle Systems* (1999).
- ▷ Funaki, *An. de l'IHP - Prob. et Stat.* (1999).
- ▷ Rezakhanlou, *Com. in math. phys.* (1991).